§ 522.88

- (c) Conditions of use in horses—(1) Amount. Administer 7 mg by intralesional injection every other day for five treatments beginning about 30 days after initial injury.
- (2) Indications for use in horses. For treatment of tendinitis of the superficial digital flexor tendon (SDFT) in horses where there is sonographic evidence of fiber tearing.
- (3) *Limitations*. Do not use in horses intended for human consumption. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[79 FR 16183, Mar. 25, 2014]

§522.88 Amoxicillin.

- (a) Specifications—(1) Each vial contains 3 grams (g) of amoxicillin trihydrate. Each milliliter of constituted suspension contains 100 or 250 milligrams (mg) amoxicillin trihydrate for use as in paragraph (d)(1) of this section.
- (2) Each vial contains 25 g of amoxicillin trihydrate. Each milliliter of constituted suspension contains 250 mg amoxicillin trihydrate for use as in paragraph (d)(2) of this section.
- (b) Sponsor. See No. 054771 in §510.600(c) of this chapter.
- (c) Related tolerance. See §556.38 of this chapter.
- (d) Conditions of use—(1) Dogs and cats—(i) Amount. Administer 5 mg per pound of body weight daily for up to 5 days by intramuscular or subcutaneous injection.
- (ii) Indications for use—(A) Dogs. For treatment of infections caused by susceptible strains of organisms as follows: Respiratory infections (tonsillitis, tracheobronchitis) due to Staphylococcus aureus, Streptococcus spp., Escherichia coli, and Proteus mirabilis; genitourinary infections (cystitis) due to S. aureus, Streptococcus spp., E. coli, and P. mirabilis; gastrointestinal infections (bacterial gastroenteritis) due to S. aureus, Streptococcus spp., E. coli, and P. mirabilis; bacterial dermatitis due to S. aureus, Streptococcus spp., and P. mirabilis; soft tissue infections (abscesses, lacerations, and wounds), due to S. aureus, Streptococcus spp., E. coli, and P. mirabilis.
- (B) Cats. For treatment of infections caused by susceptible strains of organisms as follows: Upper respiratory in-

- fections due to S. aureus, Staphylococcus spp., Streptococcus spp., Haemophilus spp., E. coli, Pasteurella spp., and P. mirabilis; genitourinary infections (cystitis) due to S. aureus, Streptococcus spp., E. coli, P. mirabilis, and Corynebacterium spp.; gastrointestinal infections due to E. coli, Proteus spp., Staphylococcus spp., and Streptococcus spp.; skin and soft tissue infections (abscesses, lacerations, and wounds) due to S. aureus, Staphylococcus spp., E. coli, and Pasteurella multocida.
- (iii) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- (2) Cattle—(i) Amount. Administer 3 to 5 mg per pound of body weight daily for up to 5 days by intramuscular or subcutaneous injection.
- (ii) Indications for use. For treatment of diseases due to amoxicillin-susceptible organisms as follows: Respiratory tract infections (shipping fever, pneumonia) due to P. multocida, P. hemolytica, Haemophilus spp., Staphylococcus spp., and Streptococcus spp. and acute necrotic pododermatitis (foot rot) due to Fusobacterium necrophorum.
- (iii) Limitations. Treated animals must not be slaughtered for food during treatment and for 25 days after the last treatment. Milk from treated cows must not be used for human consumption during treatment or for 96 hours (8 milkings) after last treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[79 FR 16183, Mar. 25, 2014]

§ 522.90 Ampicillin injectable dosage forms.

[79 FR 16183, Mar. 25, 2014]

§ 522.90a Ampicillin trihydrate suspension.

- (a) Specifications. (1) Each milliliter contains ampicillin trihydrate equivalent to 200 milligrams (mg) of ampicillin.
- (2) Each milliliter contains ampicillin trihydrate equivalent to 150 mg of ampicillin.
- (b) *Sponsors*. See sponsor numbers in §510.600(c) of this chapter.